



Уральский
федеральный
университет

имени первого Президента
России Б.Н. Ельцина

Институт
естественных наук

АНОМАЛИИ И ПАТОЛОГИИ АМФИБИЙ И РЕПТИЛИЙ:

Методология, эволюционное значение,
возможность оценки здоровья среды

Материалы международной школы-конференции
Екатеринбург, 23–26 сентября 2013 г.

МИНИСТЕРСТВО ОБРАЗОВАНИЯ И НАУКИ РОССИЙСКОЙ ФЕДЕРАЦИИ
УРАЛЬСКИЙ ФЕДЕРАЛЬНЫЙ УНИВЕРСИТЕТ
ИМЕНИ ПЕРВОГО ПРЕЗИДЕНТА РОССИИ Б. Н. ЕЛЫЦИНА
ИНСТИТУТ ЕСТЕСТВЕННЫХ НАУК

АНОМАЛИИ И ПАТОЛОГИИ АМФИБИЙ И РЕПТИЛИЙ:

Методология, эволюционное значение,
возможность оценки здоровья среды

Материалы международной школы-конференции
Екатеринбург, 23–26 сентября 2013 г.

Екатеринбург
Издательство Уральского университета
2014

Mester B., Lengyel Sz., Puky M., 2013. Occurrence of amphibian deformities in the Egyek-Pusztakócs marsh and grassland system (Hortobágy) // Programme & Abstracts Book of the 17th European Congress of Herpetology. 22–27 August 2013. Veszprém, Hungary. P. 147.

Puky M., 2006. Amphibian deformity frequency and monitoring methodology in Hungary // FROGLOG. Vol. 74. P. 3–4.

Puky M., Fodor A., 2002. Occurrence of amphibian deformities along the Hungarian section of the River Danube, Tisza and Ipoly // Limnological Reports of the International Association for Danube Research. Vol. 34. P. 845–852.

Quellet M., Bonin J., Rodrigue J., Desgranges J.-L., Lair S., 1997. Hindlimb deformities (ectromely, ectrodactyly) in free living anurans from agricultural habitats // J. Wildl. Dis. Vol. 33. P. 95–104.

Székely P., Nemes Sz., 2003. The incidence of mutilations and malformations in a population of *Pelobates fuscus* // Russ. J. of Herpetology. Vol. 10, Nr 2. P. 145–148.

Tóth M., Puky M., 2012. Herpetológiai adatgyűjtés önkéntesek fotói alapján a Kárpát-medencében: előzetes eredmények // Természetvédelmi Közlemények. Vol. 18. P. 499–505.

Vallisneri A., 1733. Opere fisico-mediche. Venezia (Venice). 551 p.

Vershinin V. L., 1989. Morphological anomalies in urban amphibians // Ékologiya. Vol. 3. P. 58–66.

TAIL LOSS AND ANOMALY IN *ZOOTOCA VIVIPARA* AND *LACERTA AGILIS* IN HUNGARY

M. Puky¹, S. Z. Faggyas², B. Mester³, P. Bíró⁴, É. Ács¹

¹ MTA Centre for Ecological Research, Danube Research Institute
(Göd, Hungary)

² Kiskunság National Park Directorate (Kecskemét, Hungary)

³ University of Debrecen, Department of Ecology (Debrecen, Hungary)

⁴ MTA Centre for Ecological Research, Balaton Limnological Institute
(Tihany, Hungary)

*An article concerned with tail's morphological anomaly that was founded between *L. agilis* from different regions of Hungary. No anomalies in *Z. vivipara* sampled were found.*

*Статья посвящена морфологической аномалии хвоста, обнаруженной среди выборок *L. agilis* из разных регионов Венгрии. Ни одной аномалии среди выборок *Z. vivipara* не отмечено.*

The study and conservation of ecological, morphological and genetic diversity of key organisms both in biodiversity hotspots and human influenced landscape is an important task at the beginning of the 21st century. The Eurasian common lizard, *Zootoca vivipara* (Lichtenstein, 1823), is the lizard with the largest distribution area on Earth [Schmidtler, Böhme, 2011] with several subspecies/clades, some with extremely large, others with more limited distribution areas. Its investigation dates back to the 1920s in Hungary [Fejérváry, 1923], where several *Z. vivipara* clades exist in different parts of the country. The study of their karyology and phylogenetic relationships started in the 2000s [Odierna et al., 2004; Surget-Groba et al., 2006] together with its conservation implications [Puky et al., 2004], which is supplemented by further and more detailed genetic, morphological and ecological investigations in recent years. As part of the study, the health status of the investigated individuals is checked. Besides *Zootoca vivipara*, *Lacerta agilis* inhabiting the same sites were also investigated morphologically. This article summarises results related with tail regeneration and anomalies.

In 2011–2013 *Zootoca vivipara* and *Lacerta agilis* was sampled in different regions of Hungary. Samples were collected from eight locations. Over a hundred individuals were examined. Individuals with additive anomalies were X-rayed.

Reptile anomalies are known for long especially that of the tail as many lizards are able to voluntarily shed their tails as a strategy to escape predation including both species studied and this process can lead to different anomalies. In the present study tail loss was recorded with a frequency over 26 % with *Z. vivipara* but no anomalies were found. A very similar frequency was found with *L. agilis*. One *Lacerta agilis* individual also showed a tail anomaly (figure). It was an adult female with a 6,83 cm body and 5,85 cm tail length and a 7,15 g weight. In the middle section of the tail (2,64 cm from its base) a 0,54 cm outgrowth was present. No bony structure was found in the outgrowth. On the basis

of digital X-ray investigations it may be related to a previous shed of the tail and its improper regeneration.



Lacerta agilis with an improper regeneration of the tail from Ökördi-láp,
Kiskőrös, Hungary (Photo: Miklós Puky)

Acknowledgements

This study was supported by the Hungarian National Research Fund (OTKA-NKTH CNK 80140). We thank Zsuzsanna Horváth, Tímea Mechura, Mónika Szalai, Sándor Kéthelyi-Nagy, András Máté, Mihály Tóth, Balázs Velekei for their participation in the fieldwork, Simon Ízing and Sándor Tacsi for X-raying and explaining the veterinary background of the phenomenon discussed above.

References

- Fejérváry G. J., 1923. Note préliminaire sur le lézard vivipare (*Lacerta vivipara* Jacq.) de la Grande Plaine Hongroise // Ann. Mus. Nat. Hung. Vol. 20. C. 166–171.
- Odierna G., Aprea G., Capriglione T., Puky M., 2004. Chromosomal evidence for the double origin of viviparity in the European common lizard, *Lacerta (Zootoca) vivipara* // Herpetological J. Vol. 14. P. 157–160.
- Puky M., Hajdu Á., Surget-Grouba Y., Heulin B., Odierna G., 2004. Fajvédelmi programok létfogosultsága és feladatai Magyarországon: az elevenszülő gyík (*Zootoca vivipara* Mayer & Bischoff, 1996) vizsgálatának eredményei és tanulságai // Természetvédelmi Közlemények. Vol. 11. P. 411–418.
- Schmidtler J. F., Böhme W., 2011. Synonymy and nomenclatural history of the Common or Viviparous Lizard, by this time: *Zootoca vivipara* (Lichtenstein, 1823) // Bonn Zoological Bulletin. Vol. 60(2). P. 214–228.
- Surget-Groba Y., Heulin B., Guillaume C.-P., Puky M., Semenov D., Orlova V., Kupriyanova L., Ghira I., Smajda B., 2006. Multiple origins of viviparity or reversal from viviparity to oviparity? The European common lizard (*Zootoca vivipara*, Lacertidae) and the evolution of parity // Biological J. of the Linnean Society. Vol. 87. P. 1–11.

**ВСТРЕЧАЕМОСТЬ МОРФОЛОГИЧЕСКИХ АНОМАЛИЙ
В ПОПУЛЯЦИОННЫХ СИСТЕМАХ
ЗЕЛЕНЫХ ЛЯГУШЕК (*PELOPHYLAX FITZINGERI*, 1843)
С СЕВЕРО-ВОСТОКА АРЕАЛОВ**

А. О. Свинин

Казанский (Приволжский) федеральный университет

**THE OCCURRENCE OF MORPHOLOGICAL ANOMALIES
IN GREEN FROGS POPULATION SYSTEMS
(*PELOPHYLAX FITZINGERI*, 1843)**

FROM THE NORTH-EASTERN PART OF THE AREAS

A. O. Svinin

Kazan (Volga Region) Federal University

In 2007–2013, a total of 754 specimens were investigated from 15 localities situated in Mari El Republic and southern part